



Title	First record of the genus <i>Sphenopus</i> (Anthozoa: Hexacorallia: Zoantharia) from Japan
Author(s)	Reimer, James Davis; Higashi, Takuo; Nonaka, Masanori
Citation	Fauna Ryukyuana, 29: 37-40
Issue Date	2016-04-19
URL	http://hdl.handle.net/20.500.12000/38739
Rights	



First record of the genus *Sphenopus* (Anthozoa: Hexacorallia: Zoantharia) from Japan

James D. Reimer^{1,2,4}, Takuo Higashiji³, Masanori Nonaka³

¹Molecular Invertebrate Systematics and Ecology Laboratory,
Department of Chemistry, Biology, and Marine Science, Faculty of Science,
University of the Ryukyus, Senbaru 1, Nishihara, Okinawa 903-0213, Japan

²Tropical Biosphere Research Center, University of the Ryukyus,
Senbaru 1, Nishihara, Okinawa 903-0213, Japan

³Okinawa Churaumi Aquarium, Okinawa Churashima Foundation,
424 Ishikawa, Motobu, Okinawa 905-0206, Japan

⁴Corresponding author (E-mail: jreimer@sci.u-ryukyu.ac.jp)

Abstract. A single specimen of *Sphenopus marsupialis* (Gmelin, 1791) was discovered within a reserve aquarium at the Okinawa Churaumi Aquarium in February 2016. Although the exact collection locality is not known the specimen clearly came from a sandy area of shallow coral reef waters around Motobu, Okinawa, Japan. Previously reported from the Central Indo-Pacific up to Taiwan, this specimen represents the first recorded occurrence of *Sphenopus* from Japanese waters based on a specimen with certain identification, as well as the northernmost record of the genus.

Introduction

The genus *Sphenopus* is distinct among zoantharians. Placed within the family Sphenopidae along with the genus *Palythoa*, *Sphenopus* species are morphologically distinct from all other genera and species as they are unitary (=monostomatous, solitary, not colonial or modular unless budding) and generally not attached to any substrate. In this regard, they are similar to the scleractinian mushroom corals (family Fungiidae), and may also have limited mobility (Soong et al. 1999). Most *Sphenopus* have large polyps of up to 3 cm in diameter and ~5 cm in length, and usually have a rounded and bulbous aboral end that is loosely anchored in sand or rubble in coral reef environments. They are azooxanthellate.

Records of *Sphenopus* spp. are not common, but they are known from the Seychelles (Den Hartog, 1997) to the central Indo-Pacific (Colin & Arneson, 1995; Erhardt & Knop, 2005; Reimer et al., 2012; 2014), and from both eastern (Zann, 1980) and western Australia (Burnett et al., 1997), with the northernmost records from Taiwan (Soong et al., 1999). Currently, there are three species within the genus. The type species *Sphenopus marsupialis* (Gmelin, 1791) is by far the most-well known, gray to tan in color with a rounded aboral end, and records

exist from a variety of locations, and there have been some studies on their reproductive ecology (Soong et al., 1999) and phylogenetic position (Reimer et al., 2012) as well. *S. arenaceus* Hertwig, 1882, is described from Cape York in northwestern Australia and before Reimer et al. (2012) had not been mentioned in scientific literature for more than 100 years. This species is morphologically similar to *S. marsupialis* but rusty red in color. The third species is *S. pedunculatus* Hertwig, 1888, from the Philippines. Unlike the other two species, this species has a 'peduncle', which is a long 'foot' or 'stalk' that is attached to small pieces of stone or rubble (see Erhardt & Knop, 2005; Reimer et al., 2014). Thus, all three species are easily distinguishable from each other and from other zoantharian species.

It is believed that *Sphenopus* species, similar to *Palythoa* species, are limited to subtropical and tropical waters. Currently, the northernmost records of *Sphenopus* are of *S. marsupialis* from Dashi, Taiwan, although this is a fishing port, and the exact locality is not exactly known (Soong et al., 1999). Additionally, there is an undocumented specimen from the East China Sea between Taiwan and the Yaeyama Islands in Uchida (2001) identified as *S. marsupialis*, but its identity is uncertain due to almost no information given on the specimen and only one small image of a single closed polyp. Despite *Sphenopus* spp.'s distinctive appearances, the relatively high number of recent investigations into the Zoantharia diversity of Okinawa (Reimer & Fujii, in press), and the relatively high Zoantharia species diversity of southern Japan (Reimer & Fujii, in press), until now there have been no records of *Sphenopus* from Japan.

Results and discussion

During a visit to Okinawa Churaumi Aquarium in




Fig. 1. Images of *Sphenopus marsupialis* specimen OCA-Cn20160220 in an aquarium at Okinawa Churaumi Aquarium, Motobu, Japan. a) Side view of specimen with polyp closed and scale (cm), b) side view at nighttime with partial opening of polyp and tentacles partially visible, and c) mostly open polyp showing two rows of tentacles.

図 1. 沖縄美ら海水族館（沖縄県本部町）の水槽内におけるダルマスナギンチャク *Sphenopus marsupialis* の生体標本（OCA-Cn20160220）。a) ポリプを閉じた側面（スケールは cm）、b) 夜間ややポリプを開いた側面、c) 二列の触手が確認できるほぼ完全に開いた状態。

Motobu, Okinawa, Japan on February 3, 2016, a single large zoantharian polyp was noticed in a reserve (=non-display) aquarium tank by the first author. The polyp was tan in color, with a rugged exterior (=tubercles), and a rounded aboral end (Fig. 1a). The polyp was approximately 5 cm in diameter and almost 9 cm in length, and was closed, with no tentacles visible (Fig. 1a). Subsequent attempts to capture images at nighttime of open polyps often failed, with a partially open oral disk with tentacles partially extended usually being observed (Fig. 1b), and only occasionally was a generally open polyp observed (Fig. 1c). The specimen is currently being housed alive in the same aquarium at Okinawa Churaumi Aquarium, and has been given specimen number OCA-Cn20160220.

Based on the polyp having a rounded aboral end, its coloration and large size, and unitary (non-colonial) morphology, the specimen was identified as *S. marsupialis* (Japanese name: darumasunaginichaku; Uchida, 2001). A subsequent search of collections records showed that the specimen was collected from waters nearby Motobu, Okinawa-jima Main Island (approximately 26° 42' N, 127° 52' E), although the exact location and depth are not known. *Sphenopus marsupialis* and other *Sphenopus* are known to inhabit sandy environments, and this specimen appears to be no exception, as it was

collected via dredging from a sandy area.

The discovery of *S. marsupialis* from Okinawa-jima Main Island is not surprising given the species' affinity for sandy tropical and subtropical environments, and also the distribution of other zoantharian species. Currently, at least 20 species of zoantharians are known from the Ryukyu Islands (Reimer & Fujii, in press; Kise & Reimer, 2016), and there are no known instances of formally described zoantharian species found in Taiwan that have not also been found in Okinawa or southern Japan. On the other hand, some Sphenopidae species are known to have their northern distributional limits in the Ryukyus, such as *Palythoa heliodiscus* (Ryland & Lancaster, 2003) with a northern limit of Amami Oshima Island (Reimer & Fujii, in press), *P. mizigama* Irei, Sinniger & Reimer, 2015 with a northern limit of Okinawa-jima Main Island, and *P. umbrosa* Irei, Sinniger & Reimer, 2015 with a northern limit of the Yaeyama Islands, while other species such as *P. tuberculosa* (Esper, 1805) and *P. mutuki* (Haddon & Shackleton, 1891) are found as far north as Miyake-jima Island in the Izu Islands south of Tokyo (Reimer et al., 2006). Future investigations should focus on examining sandy or sheltered bay regions for additional records of *Sphenopus*, as research attention has not been proportionally paid to these kinds of environments,

and they may harbor unrecorded and/or undescribed species from a wide variety of taxa (Obuchi et al., 2010; Fujii & Reimer, 2011).

Acknowledgements

This study was supported by a JSPS ‘Zuno Junkan’ grant to the first author. We thank Drs. Sandie and Bernie Degnan (University of Queensland) for providing the opportunity to visit Churaumi Aquarium. Dr. Sérgio Stampar and one anonymous reviewer are thanked for their helpful comments.

References

- Burnett, W.J., J.A.H. Benzie, J.A. Beardmore & J.S. Ryland, 1997. Zoanths (Anthozoa, Hexacorallia) from the Great Barrier Reef and Torres Strait, Australia: systematics, evolution and a key to species. *Coral Reefs*, 16: 55–68.
- Colin, P.L. & C. Arneson, 1995. *Tropical Pacific Invertebrates*. Coral Reef Press, Beverly Hills.
- Den Hartog, J.C., 1997. The sea anemone fauna of Indonesian coral reefs. In: Tomascik, T., A.J. Mah, A. Nontji & M.K. Moosa (eds.), *The ecology of the Indonesian seas 1*. Pp. 351–370, Periplus Editions, Singapore.
- Erhardt, H & D. Knop, 2005. *Corals. Indo-Pacific Field Guide*. Ikan, Frankfurt.
- Esper, E.J.C., 1805. *Die Pflanzenthiere in Abbildungen nach der Natur mit Farben erleuchtet nebst Beschreibungen*. Raspe, Nürnberg. Theilen 1–3, Lieferungen 13. [in German and Latin]
- Fujii, T. & J.D. Reimer, 2011. Phylogeny of the highly divergent family Microzoanthidae (Anthozoa, Hexacorallia) from the Pacific. *Zoologica Scripta*, 40: 418–431.
- Gmelin, J.F., 1791. *Caroli a Linne Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decimal tertia, aucta, reformata*. Georg Emanuel Beer, Lipsiae.
- Haddon, A.C. & A.M. Shackleton, 1891. Reports on the zoological collections made in Torres Straits by Professor A.C. Haddon, 1888–1889. Actiniae: I. Zoantheae. *Scientific Transactions of the Royal Dublin Society*, 4: 673–701.
- Hertwig, R., 1882. Report on the Actiniaria dredged by HMS Challenger during the years 1873–1876. Report on the scientific results of the exploring voyage of HMS Challenger 1873–1876. *Zoology*, 6: 1–122.
- Hertwig, R., 1888. Report on the Actiniaria dredged by HMS Challenger during the years 1873–1876. Supplement. Report on the scientific results of the exploring voyage of HMS Challenger 1873–1876. *Zoology*, 26: 4–56.
- Irei, Y., F. Sinniger & J.D. Reimer, 2015. Descriptions of two azooxanthellate *Palythoa* (Order Zoantharia: Subclass Hexacorallia) species from the Ryukyu Archipelago, southern Japan. *ZooKeys*, 478: 1–26.
- Kise, H. & J.D. Reimer, 2016. Unexpected diversity and new species of *Epizoanthus* (Anthozoa: Hexacorallia) attached to eunicid worm tubes from the Pacific Ocean. *ZooKeys*, 562: 49–71.
- Obuchi, M., Y. Fujita, Y. Nakano, T. Uehara & T. Motokawa, 2010. Reproductive biology and early life history of the hermaphroditic feather star *Dorometra sesokonis* (Echinodermata: Crinoidea). *Marine Biology*, 157: 1191–1201.
- Reimer, J.D. & T. Fujii, in press. Zoantharia (Cnidaria: Anthozoa: Hexacorallia) diversity research in Japan: current state and future trends. In: Kajihara, H. & Motokawa, M. (Eds.), *Species Diversity of Animals in Japan*. Springer, Tokyo.
- Reimer, J.D., A. Poliseno & B.W. Hoeksema, 2014. Shallow-water zoantherians (Cnidaria, Hexacorallia) from the Central Indo-Pacific. *ZooKeys*, 444: 1–57.
- Reimer, J.D., M. Lin, T. Fujii, D.J.W. Lane & B.W. Hoeksema, 2012. The phylogenetic position of the solitary zoanthid genus *Sphenopus* (Cnidaria: Hexacorallia). *Contributions to Zoology*, 81: 43–54.
- Reimer, J.D., S. Ono, K. Takishita, J. Tsukahara & T. Maruyama, 2006. Molecular evidence suggesting species in the zoanthid genera *Palythoa* and *Protopalythoa* (Anthozoa: Hexacorallia) are congeneric. *Zoological Science*, 23: 87–94.
- Ryland, J.S. & J.E. Lancaster, 2003. Revision for methods separating species of *Protopalythoa* (Hexacorallia: Zoanthidea) in the tropical west Pacific. *Invertebrate Systematics*, 17: 407–428.
- Soong, K., Y.S. Shiau & C.P. Chen, 1999. Morphological and life history divergence of the zoanthid, *Sphenopus marsupialis* off the Taiwanese coast. *Zoological Studies*, 38: 333–343.
- Uchida, H., 2001. *Sea Anemones in Japanese Waters*. Hankyu Communications, Tokyo. (in Japanese).
- Zann, L.P., 1980. *Living Together in the Sea*. TFH Publications, Neptune (N.J.).

ダルマスナギンチャク属（花虫綱：六放サン
ゴ亜綱：スナギンチャク目）の日本初記録

ジェイムズ D. ライマー^{1,2,4}・東地拓生³・
野中正法³

¹〒903-0213 沖縄県中頭郡西原町千原 1 番地 琉
球大学理学部海洋自然科学科生物系

²〒903-0213 沖縄県中頭郡西原町千原 1 番地 琉
球大学亜熱帯生物圏研究センター

³〒905-0206 沖縄県本部町石川 888 沖縄美ら海
水族館・(一財) 沖縄美ら島財団

要旨. 2016 年 2 月, 沖縄美ら海水族館の予備水
槽にてダルマスナギンチャク *Sphenopus*
marsupialis (Gmelin, 1791) の生体が確認された.
当該標本は沖縄県本部町周辺の浅い砂礫底で
採集されたが, その詳細な位置は定かではない.
ダルマスナギンチャク属はこれまでに中央イ
ンド太平洋から台湾まで採集の記録があるが,
日本より標本を基にして正確に同定されたの
は, 本報が初めてであり, かつ属の北限の記録
である.

投稿日: 2016 年 3 月 8 日

受理日: 2016 年 3 月 31 日

発行日: 2016 年 4 月 19 日