



Title	Effect of bioturbation activities by the grapsid crab (<i>Helice formosensis</i>) on the lipid contents and microbial consortia in tidal flat sediments: fatty acids classes-sources and fate
Author(s)	Mchenga, Islam S.S.; Tsuchiya, Makoto
Citation	琉球大学21世紀プログラム「サンゴ礁島嶼系の生物多様性の総合解析」平成17年度成果発表会
Issue Date	2006-03-06
URL	http://hdl.handle.net/20.500.12000/847
Rights	

PE-18 : Effect of bioturbation activities by the grapsid crab (*Helice formosensis*) on the lipid contents and microbial consortia in tidal flat sediments: fatty acids classes-sources and fate

Islam S.S. Mchenga, Makoto Tsuchiya

Laboratory of Ecology and Systematic, Faculty of Science, University of the Ryukyus, Senbaru 1, Nishihara, Okinawa 903 – 0213, Japan

Abstract

Macro invertebrates directly mediate key processes in the biogeochemical cycle of sediment, by releasing nutrients and consuming oxygen and organic matter. They indirectly mediate processes by redistributing particulate and dissolved substances in the sediment. A field study was conducted to investigate how burrowing activities by the grapsid crab (*Helice formosensis*) affects lipids content, fatty acid composition, and bacterial abundance in tidal flat sediments. Reported findings have shown a significant increase in total lipids in sediments of the crabs' burrow opening shaft and burrow chamber (1567.8 and 5108.2 $\mu\text{g g}^{-1}$ DW respectively, ANOVA, $p < 0.03$). This may be attributed to growth of bacteria, fungi and the product of their metabolic activities. The estimation of total microbial abundance measured by the sum of the extracted microbial derived fatty acids (bacteria and fungi biomarkers), resulted in a higher total abundance in the sediments of burrow shaft and chamber (45.7 and 24.2 $\mu\text{g g}^{-1}$ DW respectively) as compared to the surface sediments (19.9 $\mu\text{g g}^{-1}$ DW). Multivariate analyses based on principal component analysis (PCA) showed a clear separation of fatty acid methyl ester (FAMES) patterns, indicating change in bacteria makers' odd-branch fatty acids C_{15} - C_{17} (common in gram-positive bacteria) and cis-vaccenic acid $\text{C}_{18:1n-7}$ (fatty acid evidence in sulfate or sulfur reducing bacteria). These results suggest *Helice formosensis* is an active bioturbator that significantly contributes to lipid contents in biogenic organic matter by enhancing microbial abundance in the environments they inhabit.